

University of Dundee

Species coexistence in self-organised patterned ecosystems: insights from mathematical modelling


Maths seminar - 03/05/2021

Lukas Eigentler

Patterned ecosystems

- **Scale dependent feedback loops cause pattern formation** in ecological systems.
- **Local facilitation**: e.g. increased water infiltration in vegetated areas, ...
- **Long-range competition**: e.g. competition for a limiting resource.
- **Self-organisation** into colonised and uncolonised areas is typically **associated with high environmental stress**.
- Unidirectional resource flux leads to stripe patterns.

Vegetation pattern & mussel beds.




Ribbon forest




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.




Local facilitation in vegetation patterns

Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.



Local facilitation in vegetation patterns


Positive feedback loop: Water infiltration into the soil depends on local plant density \Rightarrow redistribution of water towards existing plant patches \Rightarrow further plant growth.



Patterned ecosystems

- **Scale dependent feedback loops cause pattern formation** in ecological systems.
- **Local facilitation**: e.g. increased water infiltration in vegetated areas, ...
- **Long-range competition**: e.g. competition for a limiting resource.
- **Self-organisation** into colonised and uncolonised areas is typically **associated with high environmental stress**.
- Unidirectional resource flux leads to stripe patterns.

Vegetation pattern & mussel beds.




Ribbon forest




Patterned ecosystems

- **Coexistence** typically occurs **despite competition for a single limiting resource**.
- Coexistence occurs on the scale of a single stripe (i.e. no spatial segregation).
- **What mechanisms cause coexistence?**
- Classical result: **intraspecific competition**.
- More recent: **spatial self-organisation** (i.e. local facilitation)
- What is the impact of these two contrasting processes on coexistence?

Vegetation pattern & mussel beds.



Ribbon forest



Klausmeier model

One of the most basic phenomenological models is the **extended Klausmeier reaction-advection-diffusion model**.¹

$$\begin{aligned} \frac{\partial u}{\partial t} &= \underbrace{u^2 w}_{\text{consumer growth}} - \underbrace{Bu}_{\text{consumer death}} + \underbrace{\frac{\partial^2 u}{\partial x^2}}_{\text{consumer dispersal}}, \\ \frac{\partial w}{\partial t} &= \underbrace{A}_{\text{resource input}} - \underbrace{w}_{\text{natural resource depletion}} - \underbrace{u^2 w}_{\text{resource consumption by consumers}} + \underbrace{\nu \frac{\partial w}{\partial x}}_{\text{unidirectional resource flux}} + \underbrace{d \frac{\partial^2 w}{\partial x^2}}_{\text{resource diffusion}}. \end{aligned}$$

The model describes interactions between the limiting resource and **a single consumer species**.

¹Klausmeier, C. A.: *Science* 284.5421 (1999).

Multispecies Model

Multispecies model based on the extended Klausmeier model².

$$\begin{aligned}
 \frac{\partial u_1}{\partial t} &= \underbrace{wu_1(u_1 + Hu_2)}_{\text{consumer growth}} - \underbrace{B_1 u_1}_{\text{consumer mortality}} + \underbrace{\frac{\partial^2 u_1}{\partial x^2}}_{\text{consumer dispersal}}, \\
 \frac{\partial u_2}{\partial t} &= \underbrace{Fwu_2(u_1 + Hu_2)}_{\text{consumer growth}} - \underbrace{B_2 u_2}_{\text{consumer mortality}} + \underbrace{D \frac{\partial^2 u_2}{\partial x^2}}_{\text{consumer dispersal}}, \\
 \frac{\partial w}{\partial t} &= \underbrace{A}_{\text{resource input}} - \underbrace{w}_{\text{natural resource depletion}} - \underbrace{w(u_1 + u_2)(u_1 + Hu_2)}_{\text{resource consumption by consumers}} + \underbrace{\nu \frac{\partial w}{\partial x}}_{\text{unidirectional resource flux}} + \underbrace{d \frac{\partial^2 w}{\partial x^2}}_{\text{resource diffusion}}.
 \end{aligned}$$

Species only differ quantitatively (i.e. in parameter values) but not qualitatively (i.e. same functional responses). Assume u_1 is superior coloniser; u_2 is locally superior.

²Klausmeier, C. A.: *Science* 284.5421 (1999).


Multispecies Model

Intraspecific competition (other than that for the resource) may be considered.

$$\begin{aligned}
 \frac{\partial u_1}{\partial t} &= \underbrace{wu_1(u_1 + Hu_2) \left(1 - \frac{u_1}{k_1}\right)}_{\text{consumer growth}} - \underbrace{B_1 u_1}_{\text{consumer mortality}} + \underbrace{\frac{\partial^2 u_1}{\partial x^2}}_{\text{consumer dispersal}}, \\
 \frac{\partial u_2}{\partial t} &= Fwu_2(u_1 + Hu_2) \left(1 - \frac{u_2}{k_2}\right) - \underbrace{B_2 u_2}_{\text{consumer mortality}} + \underbrace{D \frac{\partial^2 u_2}{\partial x^2}}_{\text{consumer dispersal}}, \\
 \frac{\partial w}{\partial t} &= \underbrace{A}_{\text{resource input}} - \underbrace{w}_{\text{natural resource depletion}} - \underbrace{w(u_1 + u_2)(u_1 + Hu_2)}_{\text{resource consumption by consumer}} + \underbrace{\nu \frac{\partial w}{\partial x}}_{\text{unidirectional resource flux}} + \underbrace{d \frac{\partial^2 w}{\partial x^2}}_{\text{resource diffusion}}.
 \end{aligned}$$


Species only differ quantitatively (i.e. in parameter values) but not qualitatively (i.e. same functional responses). Assume u_1 is superior coloniser; u_2 is locally superior.

Simulations





- In the **absence of intraspecific competition**, consumer coexistence is captured, but not under high environmental stress.
- Solutions are periodic travelling waves and move in the direction opposite to the unidirectional resource flux.

Simulations



- If **intraspecific competition dynamics are strong**, consumer coexistence is captured for high environmental stress.
- Solutions are periodic travelling waves and move in the direction opposite to the unidirectional resource flux.


Bifurcation diagram




- The bifurcation structure of single-species states is identical with extended Klausmeier model.

Bifurcation diagram: one wavespeed only

Bifurcation diagram




Bifurcation diagram: one wavespeed only




- The bifurcation structure of single-species states is identical with extended Klausmeier model.
- **Coexistence pattern** solution branch connects **single-species pattern** solution branches.

Pattern onset



Essential spectrum in
single-species model




Essential spectrum in
multispecies model


- The key to understand **coexistence pattern onset** is knowledge of **single-species pattern's stability**.
- Tool: **essential spectra** of periodic travelling waves, calculated using the numerical continuation method by Rademacher et al.³
- **Pattern onset occurs as the single-species pattern loses/gains stability to the introduction of a competitor.**

³Rademacher, J. D., Sandstede, B. and Scheel, A.: *Physica D* 229.2 (2007)

Pattern existence




$$B_2 - FB_1 < 0, F < 1, D < 1$$




- Key quantity: **Local average fitness difference $B_2 - FB_1$** determines stability of single-species states in spatially uniform setting.
- Condition for pattern existence: **Balance between local competitive and colonisation abilities.**

Pattern existence




$$B_2 - FB_1 \approx 0, F < 1, D < 1$$




- Key quantity: **Local average fitness difference $B_2 - FB_1$** determines stability of single-species states in spatially uniform setting.
- Condition for pattern existence: **Balance between local competitive and colonisation abilities.**

Pattern existence




$$B_2 - FB_1 > 0, F < 1, D < 1$$



- Key quantity: **Local average fitness difference $B_2 - FB_1$** determines stability of single-species states in spatially uniform setting.
- Condition for pattern existence: **Balance between local competitive and colonisation abilities.**


Pattern stability



Stability regions of system states.

- For decreasing resource input, coexistence state loses stability to single-species pattern of coloniser species.
- Transition occurs at moderate environmental stress \Rightarrow **no coexistence in the sense of a patterned ecosystem.**


Effects of intraspecific competition



Stability regions of system states.

- Intraspecific competition among colonisers stabilise coexistence in patterned state.
- Intraspecific competition among locally superior species enables spatially uniform coexistence (not shown).
- Omission of intraspecific competition leads to overestimation of single-species pattern resilience.








Hysteresis




Wavelength contours of stable patterns

- State transitions are affected by **hysteresis**.
- Extinction can occur despite a coexistence state being stable.
- **Ecosystem resilience depends on both current and past states of the system.**

Hysteresis



Spatial species distribution



- The model captures the **spatial species distribution** of grasses and trees in vegetation patterns.
- The faster the coloniser's dispersal, the more pronounced is its presence at the top edge of each stripe.

Conclusions

- **Spatial self-organisation is a coexistence mechanism.**
- Coexistence is enabled by spatial heterogeneities in the resource, caused by the consumers' self-organisation into patterns.
- A balance between species' colonisation abilities and local competitiveness promotes enables coexistence.
- **Intraspecific competition among the superior coloniser stabilises coexistence under severe environmental stress.**
- Coexistence may occur as a **metastable state** if the average fitness difference between species is small⁴.

⁴EL and Sherratt, J. A.: *Bull. Math. Biol.* 81.7 (2019).

Future Work

- How does nonlocal consumer dispersal affect species coexistence?⁵
- Do results extend to an arbitrary number of species?
- How do fluctuations in environmental conditions (in particular resource input) affect coexistence?
- In particular, what are the effects of seasonal⁶, intermittent⁷ and probabilistic resource input regimes on both single-species and multispecies states?

⁵EL and Sherratt, J. A.: *J. Math. Biol.* 77.3 (2018).




⁶EL and Sherratt, J. A.: *J. Math. Biol.* 81 (2020).

⁷EL and Sherratt, J. A.: *Physica D* 405 (2020).

References

Slides are available on my website.

<https://lukaseigentler.github.io>

-  **Eigentler, L.:** 'Species coexistence in resource-limited patterned ecosystems is facilitated by the interplay of spatial self-organisation and intraspecific competition'. *Oikos* 130.4 (2021), pp. 609–623.
-  **Eigentler, L.:** 'Intraspecific competition in models for vegetation patterns: decrease in resilience to aridity and facilitation of species coexistence'. *Ecol. Complexity* 42 (2020), p. 100835.
-  **Eigentler, L. and Sherratt, J. A.:** 'Spatial self-organisation enables species coexistence in a model for savanna ecosystems'. *J. Theor. Biol.* 487 (2020), p. 110122.